

BRIEF COMMUNICATION**Is the tiger shark *Galeocerdo cuvier* a coastal species?
Expanding its distribution range in the Atlantic Ocean using
at-sea observer data**

A. DOMINGO*†, R. COELHO‡, E. CORTES§, B. GARCIA-CORTES||, F. MAS*¶, J.
MEJUTO||, P. MILLER*¶, A. RAMOS-CARTELLE||, M. N. SANTOS‡ AND
K. YOKAWA**

*Dirección Nacional de Recursos Acuáticos (DINARA), Laboratorio de Recursos Pelágicos (LaRPe), Constituyente, 1497 Montevideo, Uruguay, ‡Portuguese Institute for the Ocean and Atmosphere (IPMA, I.P.), Avenida 5 de Outubro s/n, 8700-305, Olhão, Portugal, §National Oceanographic and Atmospheric Administration, National Marine Fisheries Service, Southeast Fisheries Science Center, Panama City Laboratory, Panama City, FL, 32408, U.S.A., ||Instituto Español de Oceanografía, P. O. Box 130, 15080, A Coruña, España, ¶Centro de Investigación y Conservación Marina (CICMAR), Av. Giannattasio Km. 30, El Pinar, Uruguay and **National Research Institute of Far Seas Fisheries, 5-7-1 Orido, Shimizu-ku, Shizuoka-City, Shizuoka, 424 8633, Japan

(Received 20 March 2015, Accepted 21 November 2015)

The occurrence of tiger shark *Galeocerdo cuvier* in the Atlantic Ocean was assessed using at-sea observer data from multiple pelagic longline fisheries. Geographic positions of 2764 *G. cuvier* recorded between 1992 and 2013 and covering a wide area of the Atlantic Ocean were compared with the currently accepted distribution ranges of the species. Most records fell outside those ranges in both the Southern and Northern Hemispheres, which strongly suggests that the distribution range of *G. cuvier* in the open ocean is considerably larger than previously described

© 2016 The Fisheries Society of the British Isles

Key words: by-catch; distribution; high seas; longline fisheries; onboard monitoring.

The tiger shark *Galeocerdo cuvier* (Peron & LeSueur 1822) is one of the most distinctive species of the family Carcharhinidae. It reaches the largest size among its congeners (up to 5.5 m total length, L_T) and is also the only carcharhinid with aplacental viviparous reproduction (Randall, 1992; Hamlett, 2005; Whitney & Crow, 2007). *Galeocerdo cuvier* is regularly described as a coastal pelagic predator with circum-global distribution in warm and temperate waters of all oceans (Ebert *et al.*, 2013). Although commonly associated with coastal areas and continental and insular shelves (Holland *et al.*, 1999; Hazin *et al.*, 2013; Afonso & Hazin, 2014), *G. cuvier* is also capable of travelling long distances, even across oceanic waters (Kohler *et al.*, 1998;

†Author to whom correspondence should be addressed. Tel.: + 598 24004689; email: adomingo@dinara.gub.uy

Heithaus *et al.*, 2007; Hammerschlag *et al.*, 2012). The distribution range of *G. cuvier* in the western Atlantic Ocean comprises coastal and shelf waters from Massachusetts, U.S.A., to Uruguay, including the Gulf of Mexico, the Caribbean and Bermuda. On the eastern Atlantic, *G. cuvier* occurs from Angola to Morocco, including the Canary and Azores archipelagos, but has also been occasionally reported close to Iceland and British Isles, possibly advected along with warm waters of the Gulf Stream (Compagno, 1984; Ebert *et al.*, 2013). To date, the presence of *G. cuvier* in the Mediterranean Sea remains uncertain and further confirmation is needed (Serena, 2005). Despite the existence of two records from Spanish and Italian waters (Pinto de la Rosa, 1994; Celona, 2000), at-sea observers of the Spanish longline fishery in the western Mediterranean Sea have not observed this species for over two decades (D. Macias, pers. comm.), suggesting that its regular occurrence in the Mediterranean Sea is doubtful.

Sub-population structure within the Atlantic Ocean is currently unknown, but long-distance migrations inferred from several tagging studies (Kohler *et al.*, 1998; Hammerschlag *et al.*, 2012) suggest that there may be connections between different regions (Mejuto *et al.*, 2005).

Galeocerdo cuvier have been not only regularly captured in some directed commercial shark fisheries (Bonfil, 1997; Simpfendorfer, 2009), recreational fisheries (Stevens, 1984) and shark control programmes (Dudley, 1997; Cliff & Dudley, 2011; Sumpton *et al.*, 2011), but also as by-catch in several fisheries (Bonfil, 1994; Beerkircher *et al.*, 2008; Coelho *et al.*, 2012). In the Atlantic Ocean, *G. cuvier* is captured as by-catch in pelagic longline fisheries across its entire distribution range. It is presently ranked as Near Threatened in the global assessment by the IUCN Red List of Threatened Species (Simpfendorfer, 2009). Although there is some information on population trends of *G. cuvier* from mostly coastal areas of the north-west Atlantic Ocean (Baum & Blanchard, 2010; Carlson *et al.*, 2012), the global trends for this species are currently unknown (Simpfendorfer, 2009). The species' conservation would be important as it is an apex predator considered as a key species in some marine ecosystems, having a considerable influence on community dynamics (Heithaus *et al.*, 2009, 2012).

Records from several observer and research programmes were pooled to assess *G. cuvier* distribution. Between 1992 and 2013, scientific observers onboard pelagic longline fishing vessels targeting tuna *Thunnus alalunga* (Bonnaterre 1788), *Thunnus obesus* (Lowe 1839), *Thunnus albacares* (Bonnaterre 1788) and *Thunnus thynnus* (L. 1758), and sword fish *Xiphias gladius* L. 1758 (Fernandez-Carvalho *et al.*, 2015) from Japan, Portugal, Spain, Uruguay and the U.S.A. recorded interactions with *G. cuvier* over a wide geographical area throughout the Atlantic Ocean, including waters over continental and insular shelves and slopes, and especially large expanses of the open ocean.

Galeocerdo cuvier were identified onboard fishing vessels by trained observers. Morphologically, *G. cuvier* is easily recognized by its short and blunt head, the presence of long upper labial furrows, low keels on the caudal peduncle, a prominent interdorsal ridge, distinctive strongly combed teeth with heavy serrations and distal cusplets, and a characteristic trunk pigmentation of dark vertical tiger-like stripe markings that become less conspicuous in adults (Compagno, 1984; Ebert *et al.*, 2013).

Data regarding the currently accepted distribution range of *G. cuvier* in the Atlantic Ocean were obtained from the IUCN Red List (IUCN, 2012) and from Ebert *et al.* (2013) (Fig. 1). Observer information collectively covered the period 1992–2013. The study was based on data for positive sets (*i.e.* fishing sets where at least one individual of

G. cuvier was captured). A buffer area of 600 km was created around the positive sets, which was then smoothed using a polynomial approximation with the exponential kernel algorithm (Peters *et al.*, 2014). Finally, the outer 200 km of the resulting polygons were clipped out in order to obtain a conservative extension of the distribution range. All spatial analysis and maps were produced using ESRI ArcGis 10.1 (www.esri.com).

In total, 2764 *G. cuvier* were reported during the study period. These were caught throughout the Atlantic Ocean, both over shelf and slope waters and their vicinity, as well as in international waters in the open ocean. Most of the records were reported by the U.S.A. (86.4%, $n = 2387$), followed by Japan (8.7%, $n = 240$), Spain (2.9%, $n = 81$), Portugal (1.3%, $n = 35$) and Uruguay (0.7%, $n = 21$).

Of all the observed *G. cuvier*, 86.4% ($n = 2388$) and 19.4% ($n = 537$) were reported outside the distribution ranges currently proposed by the IUCN (2012) and Ebert *et al.* (2013), respectively (Fig. 1). Furthermore, all *G. cuvier* reported by the Japanese, Spanish, Portuguese and Uruguayan observer programmes occurred outside the IUCN (2012) distribution ranges, as well as every individual captured over the South Atlantic Ocean. Reported catches that fell inside the currently proposed distribution ranges occurred along the U.S. east and south-east coast, the Gulf of Mexico and the Caribbean. Several reported captures also occurred over oceanic waters in areas located in the middle of the North Atlantic Ocean, over a 1000 km off the nearest coastline (Fig. 1).

Galeocerdo cuvier is infrequently captured by high-seas longline fisheries (Amorim *et al.*, 1998; Miller *et al.*, 2006; Mejuto *et al.*, 2009), probably due to its low abundance there. The observation of several hundred individuals over oceanic waters, however, confirms that this species has a more widespread distribution in the open ocean than previously accounted for. Regional satellite tagging studies in both the Atlantic (Hammerschlag *et al.*, 2012; Lea *et al.*, 2015) and Pacific Oceans (Meyer *et al.*, 2010) support the more general finding. Catches of *G. cuvier* were recorded over the entire equatorial area, whereas this does not appear to be the case for higher latitudes (Fig. 1). This suggests that the equatorial area could be acting as a connecting corridor between the east and west Atlantic, where *G. cuvier* may take advantage of both the North and South Equatorial Current and the North Equatorial Counter Current (Philander, 2001) advection as natural pathways to cross the Atlantic. Advection of *G. cuvier* by oceanic currents has also been proposed by Compagno (1984), who suggested that the isolated records reported from the U. K. could be of vagrant individuals following the Gulf and North Atlantic stream northwards. A recent telemetry study on *G. cuvier* conducted in the north-western Atlantic Ocean (Hammerschlag *et al.*, 2012) provides further evidence of long-distance migrations into the open ocean associated with waters of the Gulf Stream, and hypothesizes that these movements could be related to feeding behaviour.

Based on the data provided by several observer programmes and research activities developed on fleets that target large pelagic fishes in the high seas of the Atlantic Ocean, this paper proposes to expand the distribution range for this species (Fig. 1). This expansion is towards the open ocean, and the proposed area doubles the one previously recognized. Although the extent of observer coverage did not allow full confirmation, due to the absence of data for some fleets, it is highly possible that the apparently isolated areas appearing to the higher north and southern latitudes (question marks in Fig. 1) are also part of the distribution range, at least during certain times of the year as a consequence of migratory cycles. Expanded distribution




FIG. 1. Overlay of currently accepted distribution range of *Galeocerdo cuvier* (■, TIG range expansion IUCN, 2012) (■, Ebert *et al.*, 2013; ■, records obtained from observer and research programmes, and proposed extended distribution range in the Atlantic Ocean). ?, no positive records, but where occurrence is presumed given that there are no different conditions (★, Japan; ▲, Portugal; ■, Spain; ●, U.S.A.; ★, Uruguay).

ranges, as shown here, have direct implications for management and conservation. These findings raise questions about long-distance migrations, population or sub-population connectivity, and the identification and delimitation of different stocks. Furthermore, expanded distribution ranges also imply potentially larger availability and susceptibility to fisheries, which may result in higher vulnerability in ecological risk assessments (Cortés *et al.*, 2010). At the same time, the results suggest a potentially higher biomass than previously thought when only considering a coastal distribution. Future evaluations, such as stock and ecological risk assessments, should take this new distributional information into account to provide more realistic and up-to-date results.

This work was carried out as part of a cooperative study conducted by the ICCAT Shark Species group. The authors are grateful to all the fishery observers and longline skippers from the several nations involved in providing data for this study. Special thanks to S. Fowler and M. Dando for providing a digital version of their map. Data collection from the Portuguese fishery was funded within the scope of the EU Data Collection Framework and as part of the research projects SELECT-PAL and LL-SHARKS. R.C. was supported by a grant from the Portuguese Foundation for Science and Technology (FCT, Ref. SFRH/BPD/93936/2013). P.M. thanks the Society for Conservation GIS and the ESRI Conservation Programme for their support.

References

- Afonso, A. S. & Hazin, F. H. V. (2014). Post-release survival and behavior and exposure to fisheries in juvenile tiger sharks, *Galeocerdo cuvier*, from the South Atlantic. *Journal of Experimental Marine Biology and Ecology* **454**, 55–62.
- Amorim, A. F., Arfelli, C. A. & Fagundes, L. (1998). Pelagic elasmobranchs caught by longliners off southern Brazil during 1974–97: an overview. *Marine and Freshwater Research* **49**, 621–632.
- Baum, J. K. & Blanchard, W. (2010). Inferring shark population trends from generalized linear mixed models of pelagic longline catch and effort data. *Fisheries Research* **102**, 229–239.
- Beerkircher, L. R., Cortés, E. & Shivji, M. (2008). Case study: elasmobranch bycatch in the pelagic longline fishery off the southeastern United States, 1992–1997. In *Sharks of the Open Ocean: Biology, Fisheries and Conservation* (Camhi, M. D., Pikitch, E. K. & Babcock, E. A., eds), pp. 242–246. Oxford: Blackwell Publishing Ltd.
- Bonfil, R. (1994). Overview of world elasmobranch fisheries. *FAO Fisheries Technical Paper* **341**.
- Bonfil, R. (1997). Status of shark resources in the Southern Gulf of Mexico and Caribbean: implications for management. *Fisheries Research* **29**, 101–117.
- Carlson, J. K., Hale, L. F., Morgan, A. & Burgess, G. (2012). Relative abundance and size of coastal sharks derived from commercial shark longline catch and effort data. *Journal of Fish Biology* **80**, 1749–1764.
- Celona, A. (2000). First record of a tiger shark *Galeocerdo cuvier* (Peron & Lesueur, 1822) in the Italian waters. *Annales, Series Historia Naturalis* **10**, 207–210.
- Cliff, G. & Dudley, S. F. J. (2011). Reducing the environmental impact of shark-control programs: a case study from KwaZulu-Natal, South Africa. *Marine and Freshwater Research* **62**, 700–709.
- Coelho, R., Fernandez-Carvalho, J., Lino, P. G. & Santos, M. N. (2012). An overview of the hooking mortality of elasmobranchs caught in a swordfish pelagic longline fishery in the Atlantic Ocean. *Aquatic Living Resources* **25**, 311–319.
- Compagno, L. J. V. (1984). *FAO Species Catalogue*, Vol. 4. Sharks of the world. An annotated and illustrated catalogue of shark species known to date. Part 2. Carcharhiniformes. Rome: FAO.
- Cortés, E., Arocha, F., Beerkircher, L., Carvalho, F., Domingo, A., Heupel, M., Holtzhausen, H., Neves, M., Ribera, M. & Simpfendorfer, C. (2010). Ecological risk assessment of pelagic sharks caught in Atlantic pelagic longline fisheries. *Aquatic Living Resources* **23**, 25–34.
- Dudley, S. F. J. (1997). A comparison of the shark control programs of New South Wales and Queensland (Australia) and KwaZulu-Natal (South Africa). *Ocean and Coastal Management* **34**, 1–27.
- Ebert, D. A., Fowler, S., Compagno, L. & Dando, M. (2013). *Sharks of the World, a Fully Illustrated Guide*. Plymouth: Wild Nature Press.
- Fernandez-Carvalho, J., Coelho, R., Mejuto, J., Cortés, E., Domingo, A., Yokawa, K., Liu, K. M., García-Cortés, B., Forselledo, R., Ohshimo, S., Ramos-Cardelle, A. M., Tsai, W. P. & Santos, M. N. (2015). Pan-Atlantic distribution patterns and reproductive biology of the bigeye thresher, *Alopias superciliosus*. *Reviews in Fish Biology and Fisheries* **25**, 551–568.
- Hamlett, W. C. (2005). *Reproductive Biology and Phylogeny of Chondrichthyes: Sharks, Batoids, and Chimaeras*. Enfield, NH: Science Publishers.
- Hammerschlag, N., Gallagher, A. J., Wester, J., Luo, J. & Ault, J. S. (2012). Don't bite the hand that feeds: assessing ecological impacts of provisioning ecotourism on an apex marine predator. *Functional Ecology* **26**, 567–576.
- Hazin, F. H. V., Afonso, A. S., De Castilho, P. C., Ferreira, L. C. & Rocha, B. C. L. M. (2013). Regional movements of the tiger shark, *Galeocerdo cuvier*, off northeastern Brazil: inferences regarding shark attack hazard. *Anais da Academia Brasileira de Ciências* **85**, 1053–1062.
- Heithaus, M. R., Wirsing, A. J., Dill, L. M. & Heithaus, L. I. (2007). Long-term movements of tiger sharks satellite-tagged in Shark Bay, Western Australia. *Marine Biology* **151**, 1455–1461.

- Heithaus, M. R., Wirsing, A. J., Burkholder, D., Thomson, J. & Dill, L. M. (2009). Towards a predictive framework for predator risk effects: the interaction of landscape features and prey escape tactics. *Journal of Animal Ecology* **78**, 556–562.
- Heithaus, M. R., Wirsing, A. J. & Dill, L. M. (2012). The ecological importance of intact top-predator populations: a synthesis of 15 years of research in a seagrass ecosystem. *Marine and Freshwater Research* **63**, 1039–1050.
- Holland, K. N., Wetherbee, B. M., Lowe, C. G. & Meyer, C. G. (1999). Movements of tiger sharks (*Galeocerdo cuvier*) in coastal Hawaiian waters. *Marine Biology* **134**, 665–673.
- Kohler, N., Casey, J. & Turner, P. (1998). NMFS cooperative shark tagging program, 1962–93: an atlas of shark tag and recapture data. *Marine Fisheries Review* **60**, 1–87.
- Lea, J. S., Wetherbee, B. M., Queiroz, N., Burnie, N., Aming, C., Sousa, L. L., Mucientes, G. R., Humphries, N. E., Harvey, G. M., Sims, D. W. & Shivji, M. S. (2015). Repeated, long-distance migrations by a philopatric predator targeting highly contrasting ecosystems. *Scientific Reports* **5**, 11202. doi: 10.1038/srep11202
- Mejuto, J., García-Cortés, B. & Ramos-Cartelle, A. (2005). Tagging-recapture activities of large pelagic sharks carried out by Spain or in collaboration with the tagging programs of other countries. *ICCAT Collective Volume of Scientific Papers* **58**, 974–1000.
- Mejuto, J., García-Cortés, B., Ramos-Cartelle, A. & De la Serna, J. M. (2009). Scientific estimations of by-catch landed by the Spanish surface longline fleet targeting swordfish (*Xiphias gladius*) in the Atlantic Ocean with special reference to the years 2005 and 2006. *ICCAT Collective Volume of Scientific Papers* **64**, 2455–2468.
- Meyer, C. G., Papastamatiou, Y. P. & Holland, K. N. (2010). A multiple instrument approach to quantifying the movement patterns and habitat use of tiger (*Galeocerdo cuvier*) and Galapagos sharks (*Carcharhinus galapagensis*) at French Frigate Shoals, Hawaii. *Marine Biology* **157**, 1857–1868.
- Miller, P., Forselledo, R. & Domingo, A. (2006). Distribución de las capturas de tiburón tigre (*Galeocerdo cuvier*) por la flota Uruguaya de palangre pelágico en el Océano Atlántico Sur. In *Livro de resumos da "V Reunião da Sociedade Brasileira para o Estudo dos Elasmobrânquios (SBEEL)"* (Soto, J. M. R. & Mincarone, M. M., eds), pp. 37–38. Itajai: SBEEL/Nova Letra.
- Peters, M. P., Matthews, S. N., Iverson, L. R. & Prasad, A. M. (2014). Delineating generalized species boundaries from species distribution data and a species distribution model. *International Journal of Geographical Information Science* **28**, 1547–1560.
- Philander, S. G. (2001). Atlantic Ocean equatorial currents. In *Encyclopedia of Ocean Sciences* (Steele, J., Thorpe, S. & Turekian, K., eds), Book 1, pp. 188–191. San Diego, CA: Academic Press.
- Pinto de la Rosa, F. J. (1994). *Tiburones del mar de Alboran*. Málaga: Servicio publicaciones Centro de Ediciones de la Diputación de Málaga (CEDMA).
- Randall, J. E. (1992). Review of the biology of the tiger shark (*Galeocerdo cuvier*). *Australian Journal of Marine and Freshwater Research* **43**, 21–31.
- Serena, F. (2005). Field identification guide to the sharks and rays of the Mediterranean and Black Sea. *FAO Species Identification Guide for Fishery Purposes*. Rome: FAO.
- Stevens, J. D. (1984). Biological observations on sharks caught by sport fishermen of New South Wales. *Marine and Freshwater Research* **35**, 573–590.
- Sumpton, W. D., Taylor, S. M., Gribble, N. A., McPherson, G. & Ham, T. (2011). Gear selectivity of large-mesh nets and drumlins used to catch sharks in the Queensland Shark Control Program. *African Journal of Marine Science* **33**, 37–43.
- Whitney, N. M. & Crow, G. L. (2007). Reproductive biology of the tiger shark (*Galeocerdo cuvier*) in Hawaii. *Marine Biology* **151**, 63–70.

Electronic References

- IUCN (2012). *Galeocerdo cuvier*. In *IUCN 2012. IUCN Red List of Threatened Species. Version 2012.3*. Available at <http://www.iucnredlist.org/> (last accessed 5 May 2013).
- Simpfendorfer, C. (2009). *Galeocerdo cuvier*. In *IUCN 2013. IUCN Red List of Threatened Species, Version 2013.1*. Available at <http://www.iucnredlist.org/> (last accessed 26 August 2013).